

PARK7 single gene test

Test code: S01356

Phenotype information

- Parkinson disease, early onset

Alternative gene names

DJ-1, DJ1, GATD2

Panels that include the PARK7 gene

- [Parkinson Disease Panel](#)

Availability

3-4 weeks

Non-coding disease causing variants covered by the panel

| Gene | Genomic location HG19 | HGVS | RefSeq | RS-number |
|-------|-----------------------|-------------|-------------|-----------|
| PARK7 | Chr1:8021919 | c.-24+66C>G | NM_007262.4 | |

Test Strengths

The strengths of this test include:

- CAP accredited laboratory
- CLIA-certified personnel performing clinical testing in a CLIA-certified laboratory
- Powerful sequencing technologies, advanced target enrichment methods and precision bioinformatics pipelines ensure superior analytical performance
- Careful construction of clinically effective and scientifically justified gene panels
- Our Nucleus online portal providing transparent and easy access to quality and performance data at the patient level
- Our publicly available analytic validation demonstrating complete details of test performance
- ~2,000 non-coding disease causing variants in our clinical grade NGS assay for panels (please see 'Non-coding disease causing variants covered by this test')
- Our rigorous variant classification scheme
- Our systematic clinical interpretation workflow using proprietary software enabling accurate and traceable processing of NGS data
- Our comprehensive clinical statements

Test Limitations

This test does not detect the following:

- Complex inversions
- Gene conversions

- Balanced translocations
- Mitochondrial DNA variants
- Repeat expansion disorders unless specifically mentioned
- Non-coding variants deeper than ± 20 base pairs from exon-intron boundary unless otherwise indicated (please see above non-coding variants covered by the panel).

This test may not reliably detect the following:

- Low level mosaicism (variant with a minor allele fraction of 14.6% is detected with 90% probability)
- Stretches of mononucleotide repeats
- Indels larger than 50bp
- Single exon deletions or duplications
- Variants within pseudogene regions/duplicated segments

The sensitivity of this test may be reduced if DNA is extracted by a laboratory other than Blueprint Genetics.

For additional information, please refer to the Test performance section.

Test Performance

Our single gene tests are sectioned from our high-quality, clinical grade NGS assay. Please see our sequencing and detection performance table for details regarding our ability to detect different types of alterations (Table).

Assays have been validated for various sample types including EDTA-blood, isolated DNA (excluding from formalin fixed paraffin embedded tissue), saliva and dry blood spots (filter cards). These sample types were selected in order to maximize the likelihood for high-quality DNA yield. The diagnostic yield varies depending on the assay used, referring healthcare professional, hospital and country. Plus analysis increases the likelihood of finding a genetic diagnosis for your patient, as large deletions and duplications cannot be detected using sequence analysis alone. Blueprint Genetics' Plus Analysis is a combination of both sequencing and deletion/duplication (copy number variant (CNV)) analysis.

The performance metrics listed below are from an initial validation performed at our main laboratory in Finland. The performance metrics of our laboratory in Seattle, WA, are equivalent.

Performance of Blueprint Genetics high-quality, clinical grade NGS sequencing assay for single genes.

| | Sensitivity % (TP/(TP+FN)) | Specificity % |
|---|----------------------------|---------------|
| Single nucleotide variants | 99.89% (99,153/99,266) | >99.9999% |
| Insertions, deletions and indels by sequence analysis | | |
| 1-10 bps | 99.2% (7,745/7,806) | >99.9999% |
| 11-50 bps | 99.13% (2,524/2,546) | >99.9999% |
| Copy number variants (exon level dels/dups) | | |
| 1 exon level deletion (heterozygous) | 100% (20/20) | NA |
| 1 exon level deletion (homozygous) | 100% (5/5) | NA |
| 1 exon level deletion (het or homo) | 100% (25/25) | NA |
| 2-7 exon level deletion (het or homo) | 100% (44/44) | NA |
| 1-9 exon level duplication (het or homo) | 75% (6/8) | NA |
| Simulated CNV detection | | |



| | | |
|--|--------------|---------|
| 5 exons level deletion/duplication | 98.7% | 100.00% |
| Microdeletion/-duplication sdrs (large CNVs, n=37) | | |
| Size range (0.1-47 Mb) | 100% (25/25) | |

The performance presented above reached by Blueprint Genetics high-quality, clinical grade NGS sequencing assay with the following coverage metrics

| | |
|---|--------|
| Mean sequencing depth | 143X |
| Nucleotides with >20x sequencing coverage (%) | 99.86% |

Performance of Blueprint Genetics Mitochondrial Sequencing Assay.

| | Sensitivity % | Specificity % |
|---|--------------------|---------------|
| ANALYTIC VALIDATION (NA samples; n=4) | | |
| Single nucleotide variants | | |
| Heteroplasmic (45-100%) | 100.0% (50/50) | 100.0% |
| Heteroplasmic (35-45%) | 100.0% (87/87) | 100.0% |
| Heteroplasmic (25-35%) | 100.0% (73/73) | 100.0% |
| Heteroplasmic (15-25%) | 100.0% (77/77) | 100.0% |
| Heteroplasmic (10-15%) | 100.0% (74/74) | 100.0% |
| Heteroplasmic (5-10%) | 100.0% (3/3) | 100.0% |
| Heteroplasmic (<5%) | 50.0% (2/4) | 100.0% |
| CLINICAL VALIDATION (n=76 samples) | | |
| All types | | |
| Single nucleotide variants n=2026 SNVs | | |
| Heteroplasmic (45-100%) | 100.0% (1940/1940) | 100.0% |
| Heteroplasmic (35-45%) | 100.0% (4/4) | 100.0% |
| Heteroplasmic (25-35%) | 100.0% (3/3) | 100.0% |
| Heteroplasmic (15-25%) | 100.0% (3/3) | 100.0% |
| Heteroplasmic (10-15%) | 100.0% (9/9) | 100.0% |
| Heteroplasmic (5-10%) | 92.3% (12/13) | 99.98% |
| Heteroplasmic (<5%) | 88.9% (48/54) | 99.93% |
| Insertions and deletions by sequence analysis n=40 indels | | |



Blueprint Genetics



| | | |
|---|-----------------|-------------------|
| Heteroplasmic (45-100%) 1-10bp | 100.0% (32/32) | 100.0% |
| Heteroplasmic (5-45%) 1-10bp | 100.0% (3/3) | 100.0% |
| Heteroplasmic (<5%) 1-10bp | 100.0% (5/5) | 99,997% |
| SIMULATION DATA /(mitomap mutations) | | |
| Insertions, and deletions 1-24 bps by sequence analysis; n=17 | | |
| Homoplasmic (100%) 1-24bp | 100.0% (17/17) | 99.98% |
| Heteroplasmic (50%) | 100.0% (17/17) | 99.99% |
| Heteroplasmic (25%) | 100.0% (17/17) | 100.0% |
| Heteroplasmic (20%) | 100.0% (17/17) | 100.0% |
| Heteroplasmic (15%) | 100.0% (17/17) | 100.0% |
| Heteroplasmic (10%) | 94.1% (16/17) | 100.0% |
| Heteroplasmic (5%) | 94.1% (16/17) | 100.0% |
| Copy number variants (separate artificial mutations; n=1500) | | |
| Homoplasmic (100%) 500 bp, 1kb, 5 kb | 100.0% | 100.0% |
| Heteroplasmic (50%) 500 bp, 1kb, 5 kb | 100.0% | 100.0% |
| Heteroplasmic (30%) 500 bp, 1kb, 5 kb | 100.0% | 100.0% |
| Heteroplasmic (20%) 500 bp, 1kb, 5 kb | 99.7% | 100.0% |
| Heteroplasmic (10%) 500 bp, 1kb, 5 kb | 99.0% | 100.0% |
| The performance presented above reached by following coverage metrics at assay level (n=66) | | |
| | Mean of medians | Median of medians |
| Mean sequencing depth MQ0 (clinical) | 18224X | 17366X |
| Nucleotides with >1000x MQ0 sequencing coverage (%) (clinical) | 100% | |
| rho zero cell line (=no mtDNA), mean sequencing depth | 12X | |

Bioinformatics

The target region for each gene includes coding exons and ± 20 base pairs from the exon-intron boundary. In addition, the panel includes non-coding and regulatory variants if listed above (Non-coding variants covered by the panel). Some regions of the gene(s) may be removed from the panel if specifically mentioned in the "Test limitations" section above. The sequencing data generated in our laboratory is analyzed with our proprietary data analysis and annotation pipeline, integrating state-of-the-art algorithms and industry-standard software solutions. Incorporation of rigorous quality control steps throughout the workflow of the pipeline ensures the consistency, validity and accuracy of results. Our pipeline is streamlined to maximize sensitivity without sacrificing specificity. We have incorporated a number of reference population databases and mutation databases including, but not limited, to [1000 Genomes Project](#), [gnomAD](#), [ClinVar](#) and [HGMD](#) into our clinical interpretation software to make the process effective and efficient. For missense variants, *in silico* variant prediction tools such as [SIFT](#), [PolyPhen](#), [MutationTaster](#) are used to assist with variant classification. Through our online ordering and statement

Blueprint Genetics



reporting system, Nucleus, ordering providers have access to the details of the analysis, including patient specific sequencing metrics, a gene level coverage plot and a list of regions with <20X sequencing depth if applicable. This reflects our mission to build fully transparent diagnostics where ordering providers can easily visualize the crucial details of the analysis process.

Clinical Interpretation

We provide customers with the most comprehensive clinical report available on the market. Clinical interpretation requires a fundamental understanding of clinical genetics and genetic principles. At Blueprint Genetics, our PhD molecular geneticists, medical geneticists and clinical consultants prepare the clinical statement together by evaluating the identified variants in the context of the phenotypic information provided in the requisition form. Our goal is to provide clinically meaningful statements that are understandable for all medical professionals regardless of whether they have formal training in genetics.

Variant classification is the corner stone of clinical interpretation and resulting patient management decisions. Our classifications follow the [ACMG guideline 2015](#).

The final step in the analysis is orthogonal confirmation. Sequence variants classified as pathogenic, likely pathogenic and variants of uncertain significance (VUS) are confirmed using bi-directional Sanger sequencing when they do not meet our stringent NGS quality metrics for a true positive call. Reported heterozygous and homo/hemizygous copy number variations with a size <10 and <3 target exons are confirmed by orthogonal methods such as qPCR if the specific CNV has been seen and confirmed less than three times at Blueprint Genetics.

Our clinical statement includes tables for sequencing and copy number variants that include basic variant information (genomic coordinates, HGVS nomenclature, zygosity, allele frequencies, in silico predictions, OMIM phenotypes and classification of the variant). In addition, the statement includes detailed descriptions of the variant, gene and phenotype(s) including the role of the specific gene in human disease, the mutation profile, information about the gene's variation in population cohorts and detailed information about related phenotypes. We also provide links to the references, abstracts and variant databases used to help ordering providers further evaluate the reported findings if desired. The conclusion summarizes all of the existing information and provides our rationale for the classification of the variant.

Identification of pathogenic or likely pathogenic variants in dominant disorders or their combinations in different alleles in recessive disorders are considered molecular confirmation of the clinical diagnosis. In these cases, family member testing can be used for risk stratification. We do not recommend using variants of uncertain significance (VUS) for family member risk stratification or patient management. Genetic counseling is recommended.

Our interpretation team analyzes millions of variants from thousands of individuals with rare diseases. Our internal database and our understanding of variants and related phenotypes increases with every case analyzed. Our laboratory is therefore well-positioned to re-classify previously reported variants as new information becomes available. If a variant previously reported by Blueprint Genetics is re-classified, our laboratory will issue a follow-up statement to the original ordering healthcare provider at no additional cost, according to our latest follow-up reporting policy.

CPT code(s) *

81479

* The CPT codes provided are based on AMA guidelines and are for informational purposes only. CPT coding is the sole responsibility of the billing party. Please direct any questions regarding coding to the payer being billed.

Sample Requirements

- Blood (min. 1ml) in an EDTA tube
- Extracted DNA, min. 2 µg in TE buffer or equivalent
- Saliva (Please see [Sample Requirements](#) for accepted saliva kits)

Label the sample tube with your patient's name, date of birth and the date of sample collection.

Blueprint Genetics



We do not accept DNA samples isolated from formalin-fixed paraffin-embedded (FFPE) tissue. In addition, if the patient is affected with a hematological malignancy, DNA extracted from a non-hematological source (e.g. skin fibroblasts) is strongly recommended.

Please note that, in rare cases, mitochondrial genome (mtDNA) variants may not be detectable in blood or saliva in which case DNA extracted from post-mitotic tissue such as skeletal muscle may be a better option.

Read more about our sample requirements [here](#).